

## Case Report

# Two Cases of Peritonitis due to Pantoea Species at One Center with Different Outcomes

Essam S, Noor A, Eldin MAN, Alsaleh E, Abdurhman IS and Alhwiesh A\*

Department of Internal Medicine, Imam Abdulrahman Bin Faisal University, Saudi Arabia

\*Corresponding author: Alhwiesh A, Department of Internal Medicine, Imam Abdulrahman Bin Faisal University, Division of Nephrology, King Fahd Hospital of the University, Al-Khobar, Saudi Arabia

Received: December 11, 2017; Accepted: January 24, 2018; Published: January 31, 2018

## Abstract

Peritonitis is a serious complication of peritoneal dialysis. A variety of microorganisms are identified in these cases and during recent years a new one was included, *Pantoea* species. In our report, we present 2 cases of patients on CCPD with a peritonitis episode caused by this organism. The source of infection in one of the cases was thought to be due to gardening of the plant Basil, while unknown in the other case. In microbiologic culture, this organism was identified and the patients were started on antibiotics with success in one case while the other requiring catheter removal. The number of reported cases with this organism has increased in last years and various infection localizations and clinical progress patterns have been identified. In peritoneal dialysis patients presenting with peritonitis, this organism must be kept in mind.

**Keywords:** Peritonitis; *Pantoea* species; Peritoneal dialysis

## Introduction

For patients on peritoneal dialysis, peritonitis is an important cause of mortality and morbidity. The treating physician needs to keep a high index of suspicion and treat peritonitis early to improve outcome and reduce complications. While usual gram positive and gram negative organisms make most of the cases, recently one unusual environmental gram negative family appeared to be the cause.

The organism *Pantoea*, belongs to the family Enterobacteriaceae and is responsible for infectious diseases mainly from plant-thorn injuries, causing arthritis, osteoarthritis, osteomyelitis to bacteremia. Until now a number of clinical cases of peritonitis caused by this organism have been described.

We present two cases (Case A and B) of peritonitis with *Pantoea* species at our center, both of which had different course and outcomes requiring different treatments.

## Case Reports

### Case A

The first case was a 19 year old male who has been on Peritoneal Dialysis for last 2 years without complications and no history of previous peritonitis, presented with 3 days history of abdominal pain and fever. The symptoms started two days after spending time gardening a specific type of plant called Reyhan - (Basil) at his home, although the patient did not remember any event of thorn pricks while planting Figure 1.

Upon evaluation, there was diffuse abdominal tenderness but there were no signs of exit site or tunnel infection. The PD fluid was found to be cloudy and analysis showed White cell count of 1000 with 92% neutrophils. Other investigations revealed serum WBC count 7.6k/ul, ESR 30mm/hr, CRP 0.4mg/dl, Lactic acid 3.1mmol/L. Patient was started on empirical treatment including Intraperitoneal Ceftriaxone and Vancomycin. The Blood cultures were negative; however the PD fluid culture grew *Pantoea* species sensitive to

Cephalosporins, Gentamycin and Tazobactam. The patient showed significant improvement and disappearance of symptoms on the third day. The PD fluid WBC dropped to 22k/ul with normalization of ESR and lactic acid levels and the fluid culture became negative after 5 days of treatment. Patient was discharged home and he completed 14 days of antibiotic therapy. Over the next one year, patient did not have any further episodes of peritonitis, and the repeated PET test did not show any changes on the membrane characteristics.

### Case B

The second case was a 33 year old female who was receiving APD for 3 years and had two prior episodes of culture negative peritonitis.

She had been visiting Paris for 2 weeks but did not report any significant events such as injuries cuts or pricks. She developed vomiting, abdominal discomfort and diarrhea two days before presentation. Upon evaluation patient was ill looking with fever, and diffuse abdominal tenderness. The PD fluid was slightly hazy with WBC count of 602 and 83% neutrophils. Her Serum WBC count was 3.9k/ul, ESR 67mm/hr, CRP 8.3mg/dl, Lactic acid 1.4mmol/L. She was started empirically on Intraperitoneal Ceftriaxone and Vancomycin. The PD fluid culture was released on 4<sup>th</sup> day which grew *Pantoea* Species, sensitive to Ceftazidime, Gentamycin and Tazocin Figure 2.

She failed to show any significant clinical improvement in the following days and the repeated PD fluid white cell count continued to rise on the 3<sup>rd</sup>, 5<sup>th</sup> and 7<sup>th</sup> days. The antibiotics were changed to Meropenam and Gentamycin. Due to persistent abdominal pain, it was decided to remove the catheter on 11<sup>th</sup> day, following which remarkable improvement was noticed. She was given total 3 weeks of the last antibiotic regimen and was shifted to hemodialysis temporarily, returning back to peritoneal dialysis in about 3 months time. No significant changes were observed on subsequent PET tests.

## Discussion

Peritonitis is a serious complication of peritoneal dialysis [1] and early treatment is essential to prevent mortality and morbidity



**Figure 1:** Basil Plant from patient's garden.

including future changes in membrane characteristics and possible ultrafiltration failure. In recent years, increasing number of cases due to rather uncommon organisms has been reported. One such genus is *Pantoea*, a Gram-negative aerobic bacillus belonging to the Enterobacteriaceae family. This organism is found especially in plants, fruits, vegetables and also human and animal feces [2]. Its transmission is generally associated with plant thorn injuries, intravenous anaesthetic catheters, contaminated parenteral nutrition and subgingival sites with periodontal diseases [3,4]. Reported infections include osteoitis, osteomyelitis, arthritis [8] to systemic infections and sepsis. Peritonitis is a rare presentation of this organism. The symptoms of *Pantoea* associated peritonitis are similar to those with other pathogens, including fever, nausea, vomiting and abdominal pain. It can cause symptomatic peritonitis with a fatal outcome. Therefore, microbiological evaluation and antibiotic susceptibility tests should be required for every peritonitis episode as suggested by International Society for Peritoneal Dialysis (ISPD) guidelines [5]. *Pantoea* peritonitis shows good response to antibiotics.

We are reporting two cases of *Pantoea* peritonitis in our PD patients. One patient had probably acquired from gardening of plant Basil, while the other did not have a clear causal event. Growth of *Pantoea* in both patients was highly sensitive to most antibiotics, however one patient did not respond well to antibiotics and required removal of catheter Table 1.

**Table 1:**

| Case(Ref.) | Type pf PD | Sex/Age | Proposed Etiology    | Antibiotics used/route    | Outcome                |
|------------|------------|---------|----------------------|---------------------------|------------------------|
| 1 (12)     | NA         | F/2     | Teething of catheter | Cefotaxime, Gentamycin/IP | Cured, Cathterreplaced |
| 2 (13)     | CAPD       | F/49    | Rose-Thorn Injury    | Ceftazidime, Amikacin/IP  | Cured                  |
| 3 (6)      | CCPD       | M/65    | Unknown              | Ciprofloxacin/IP          | Cured                  |
| 4 (7)      | CAPD       | M/51    | Rose-Thorn Injury    | Cefepime/IP               | Cured                  |
| 5 (14)     | NA         | F/52    | Unknown              | Ciprofloxacin/Oral        | Cured                  |
| 6 (15)     | CAPD       | M/45    | Thorn Injury         | Ciprofloxacin/IP          | Cured                  |
| 7 (16)     | CCPD       | M/56    | Unknown              | Tobramycin                | Cured                  |
| 8 (4)      | APD        | F/89    | Unknown              | Imipenam/IV               | Patient Died           |
| Our Case A | APD        | M/19    | Thorn Injury (Basil) | Ceftriaxone/IP            | Cured                  |
| Our Case B | APD        | F/33    | Unknown              | Meropenam, Gentamycin     | Cured, Cathter removed |



**Figure 2:** *Pantoea* agglomerans strain producing yellow pigmentation on blood agar.

In review of literature, we found that only a few case reports of *Pantoea* associated peritonitis. The mode of infection and pathogenesis of this disease in this patients' group remained unclear. There is no proven better diagnostic technique for identification. The incubation period is uncertain and inappropriate growth media and identification methods could be the reason for negative results. Different contamination patterns were identified like rose-thorn injury and teething on a catheter in these reports also [12-14]. However, some patients proposed etiologies were unknown [14,15].

Response to treatment is favorable and early treatment has resulted in successful resolution of peritonitis in majority of cases. In PD patients, peritoneal function tests may be affected after this infection. Ultrafiltration failure has been observed in some cases and peritoneal calcification and thickening were considered as possible pathologies [16].

### Conclusion

In summary, relatively new organisms are recently reported to be causing peritonitis in PD patients. *Pantoea* has also been reported and has been treated successfully with antibiotics, either with or without removal of catheter, while causing one mortality. One of our presented patients responded to antibiotics alone, while the other required removal of catheter. Further studies are required to ascertain the mode of transmission and the adequate duration and modality of treatment.

### References

1. Troidle L, Gorban-Brennan N, Kliger A, Finkelstein FO. Continuous peritoneal

- dialysis-associated peritonitis: a review and current concepts. *Semin Dial.* 2003; 16: 428-437.
2. Bottone E, Schneerson SS. *Erwinia* species: an emerging human pathogen. *Am J Clin Pathol.* 1972; 57: 400-405.
  3. Kratz A, Greenberg D, Barki Y, Cohen E, Lifshitz M. Pantoea agglomerans as a cause of septic arthritis after palm tree thorn injury; case report and literature review. *Arch Dis Child.* 2003; 88: 542-544.
  4. Kahveci A, Ascioglu E, Tigen E, Ari E, Arikan H, Odabasi Z, et al. Unusual causes of peritonitis in a peritoneal dialysis patient: *Alcaligenes faecalis* and *Pantoea agglomerans*. *Ann Clin Microbiol Antimicrob.* 2011; 10: 12.
  5. Li PK, Szeto CC, Piraino B, Bernardini J, Figueiredo AE, Gupta A, et al. Peritoneal dialysis-related infections recommendations 2010 update. *Perit Dial Int.* 2010; 30: 393-423.
  6. Magnette C, Tintillier M, Horlait G, Cuvelier C, Pochet JM. Severe peritonitis due to *Pantoea agglomerans* in a CCPD patient. *Perit Dial Int.* 2008; 28: 207-208.
  7. Ferrantino M, Navaneethan SD, Sloand JA. *Pantoea agglomerans*: an unusual inciting agent in peritonitis. *Perit Dial Int.* 2008; 28:428-430.
  8. Ulloa-Gutierrez R, Moya T, Avila-Aguero ML. *Pantoea agglomerans* and thorn-associated suppurative arthritis. *Pediatr Infect Dis J.* 2004; 23:690.
  9. Lau KK, Ault BH, Jones DP. Polymicrobial peritonitis including *Pantoea agglomerans* from teething on a catheter. *South Med J.* 2005; 98: 580-581.
  10. Lim PS, Chen SL, Tsai CY, Pai MA. *Pantoea* peritonitis in a patient receiving chronic ambulatory peritoneal dialysis. *Nephrology.* 2006; 11: 97-99.
  11. Habhab W, Blake PG. *Pantoea* peritonitis: not just a "thorny" problem. *Perit Dial Int.* 2008; 28: 430.
  12. Moreiras-Plaza M, Blanco-García R, Romero-Jung P, FeijóoPiñeiro D, Fernandez-Fernandez C, Ammari I. *Pantoea agglomerans*: the gardener's peritonitis? *Clin Nephrol.* 2009; 72: 159-161.
  13. Borrás M, Roig J, Garcia M, Fernández E. Adverse effects of *pantoea* peritonitis on peritoneal transport. *Perit Dial Int.* 2009; 29: 234-235.