

## Perspective

# Transcranial Ultrasound in Neurocritical Patient (TUSiNP Protocol): A Proposal

Abdo-Cuza A\*

Centro de Investigaciones Médico Quirúrgicas, La Habana, Cuba

\*Corresponding author: Anselmo Abdo-Cuza, Centro de Investigaciones Médico Quirúrgicas, La Habana, Cuba

Received: May 08, 2019; Accepted: June 04, 2019;

Published: June 11, 2019

## Perspective

The use of multimodal monitoring in neurocritical patients meant a milestone in neurocritical care [1]. Invasive measurement of intracranial pressure is still the gold standard method for evaluating cerebral hemodynamic [2]. However, there is a group of clinical situations or care settings in which the use of non-invasive methods of neuromonitoring would be ideal.


At the end of the 80s, color-coded transcranial duplex ultrasound (CCTD) began to be introduced into the adult population, a technique that combines the acquisition of color and spectral images by transcranial Doppler together with two-dimensional images [3]. Its generalization in medical practice has not been parallel to the point of care ultrasound (POCUS). There is a group of authors who try to make the transcranial ultrasound part of POCUS.

The method requires a learning curve not different from that required by critical ultrasound and an approach based on pathophysiological knowledge of the normal and damaged brain. In POCUS the use of some basic protocols such as RUSH, FAST, e-FAST, among others has facilitated their learning and generalization [4,5].

Perhaps the proposal of a protocol for the use of transcranial ultrasound in neurocritical patient (TUSiNP) could help the implementation of the method. At this point it is mandatory to highlight its use together with clinical examination and judgment.

The same ultrasound used in the evaluation of critical patients is used, with the same sectoral transducer used for echocardiography (1-5 MHz). Through the transtemporal window the views of the diencephalic, mesencephalic and ventricular planes will be obtained. Once the third ventricle has been identified and the distance from it measured from each side, it will be possible to assess whether there is a deviation from the midline [6].

In the bi-dimensional images, the presence of intracranial hematomas can be evaluated [7]. In a transtemporal window in the mesencephalic and diencephalic planes, the blood vessels that make up the polygon of Willis can be visualized. By identifying the vessels by color Doppler and placing the pulsed Doppler sample volume in them, Doppler spectra of flow velocities are obtained (Figure 1). The evaluation of mean velocity, final diastolic velocity and pulsatility index allow an approximation to the pattern cerebral hemodynamics [6,8]. Five patterns can be diagnosed: low flow, high resistance,



**Figure 1:** Color and spectral Doppler. Insonation of middle cerebral artery by transtemporal window.

hyperemic, vasospasm and circulatory arrest. The identification allows to initiate personalized behaviors and to have differential diagnoses in each pattern. There are described formulas for the non-invasive estimation of intracranial pressure (ICP) and cerebral perfusion pressure [9]. Acceptance is controversial but trends can be followed.

With a 3-8 MHz linear transducer through the transtemporal window, the diameter of the optic nerve sheath can be determined [10]. An increase above parameters that must be established corresponds to an increase in ICP.

In summary, color-coded transcranial duplex ultrasound can have an important value for the non-invasive evaluation of cerebral hemodynamics, by following an established protocol: TUSiNP protocol.

## References

1. Makarenko S, Griesdale DE, Gooderham P, Sekhon MS. Multimodal neuromonitoring for traumatic brain injury: A shift towards individualized therapy. *J Clin Neurosci*. 2016; 26: 8-13.
2. Kawoos U, Mc Carron RM, Auker CR, Chavko M. Advances in Intracranial Pressure Monitoring and Its Significance in Managing Traumatic Brain Injury. *Int J Mol Sci*. 2015; 16: 28979-28997.
3. Blanco P. Transcranial Color-Coded Duplex Sonography: Another Option besides the Blind Method. *J Ultrasound Med*. 2016; 35: 669-671.
4. Bagheri-Hariri S, Yekesadat M, Farahmand S, Arbab M, Sedaghat M, Shahlafar N, et al. The impact of using RUSH protocol for diagnosing the type of unknown shock in the emergency department. *Emerg Radiol*. 2015; 22: 517-520.
5. Flato UA, Guimarães HP, Lopes RD, Valiatti JL, Flato EM, Lorenzo RG. Usefulness of Extended-FAST (EFAST-Extended Focused Assessment with Sonography for Trauma) in critical care setting. *Rev Bras Ter Intensiva*. 2010; 22: 291-299.

6. Blanco P, Abdo-Cuza A. Transcranial Doppler ultrasound in neurocritical care. *J Ultrasound*. 2018; 21: 1-16.
7. Blanco P, Matteoda M. Images in emergency medicine. Extra-axial intracranial hematoma, midline shift, and severe intracranial hypertension detected by transcranial color-coded duplex sonography. *Ann Emerg Med*. 2015; 65: e1-2.
8. Abdo A, Pérez-Bernal J, Hinojosa R, Porras F, Castellanos R, Gómez F, et al. Cerebral Hemodynamics Patterns by Transcranial Doppler in Patients With Acute Liver Failure. *Transplant Proc*. 2015; 47: 2647-2649.
9. Robba C, Cardim D, Tajsic T, Pietersen J, Bulman M, Donnelly J, et al. Ultrasound non-invasive measurement of intracranial pressure in neurointensive care: A prospective observational study. *PLoS Med*. 2017; 14: e1002356.
10. Koziarz A, Sne N, Kegel F, Alhazzani W, Nath S, Badhiwala JH, et al. Optic nerve sheath diameter sonography for the diagnosis of increased intracranial pressure: a systematic review and meta-analysis protocol. *BMJ Open*. 2017; 7: e016194.