

Special Article - Helicobacter Pylori Infection

Helicobacter Pylori and Nissen Funduplication

Polat FR*

Division of Surgery, NK University, Medical Faculty,
59100 Tekirdag, Turkey

***Corresponding author:** Fatin Rüştü Polat, Division
of Surgery, NK University, Medical Faculty, 59100
Tekirdag, Turkey

Received: December 17, 2016; **Accepted:** December 29,
2016; **Published:** January 02, 2017

Letter to the Editor

Most of the patients who underwent antireflux surgery; a little time after surgery many of symptom is seen relapse. this condition is not false surgery. But this is result of insufficient evaluatin preoperatively. Such as helicobacter pylori, dental plaque. Surgeon should be sure enough gastric outlet and oral hygiene. Before surgery, biopsy must be taken on antrum and check oral cavity absolutely [1,2].

As a result, Antireflux surgery candidate patients should be examined on Helicobacter pylori. Because Helicobacter pylori is accumulated in antrum. This condition is cause of gastric outlet obstruction. Dental plaque, poor oral hygiene, and periodontal disease are risk factors for Helicobacter pylori infection.

References

1. Polat FR, Polat S. The effect of Helicobacter pylori on gastroesophageal reflux disease. JLS. 2012; 16: 260- 263.
2. Carini Francesco, Leone Angelo, Di Pasquale Roberto, et all. Association Between Helycobacter Pylori Infection and Pathological Oral Manifestations. J Int Transl Med. 2016; 4: 9-15.