

Research Article

The Neurosurgical Management of Peroneal Nerve Injury: An Experience of 14 Cases

Simas RT^{1,2}, Caires ACV¹, Monteiro PIP^{1,2}, Dantas F^{1,2} and Dantas FLR^{1,3*}

¹Department of Neurosurgery, Biocor Instituto, Nova Lima, Minas Gerais, Brazil

²Post-Graduation, Faculdade de Ciências Médicas de Minas Gerais - Feluma, Belo Horizonte, Minas Gerais, Brazil

³Post-Graduation, Instituto de Assistência Médica ao Servidor Público Estadual de São Paulo - IAMSPE, São Paulo, São Paulo, Brazil

*Corresponding author: Dantas FLR, Department of Neurosurgery, Biocor Instituto, Alameda Oscar Niemeyer, 217, Nova Lima, Minas Gerais, Brazil

Received: June 14, 2021; Accepted: July 13, 2021;

Published: July 20, 2021

Abstract

Objectives: To describe the neurosurgical technical nuances of peroneal nerve injury management and to analyze the outcomes of patients diagnosed with peroneal nerve injury operated on at a single institution.

Methods: Fourteen patients, all with electroneuromyography confirmation of peroneal nerve injury, were retrospectively analyzed. The variables analyzed included patient demographic characteristics, etiology of the lesion, preoperative neurological status, location of the lesion, perioperative findings, surgical technique, complications, and neurological status six months postoperatively.

Results: Traumatic injury was the most common cause of peroneal nerve injury, accounting for 64.27% of cases. Concerning surgical technique, neurolysis was the preferred technique in most cases. Isolated neurolysis was performed in 50% of the cases, neurolysis combined with graft in 7.14%, and neurolysis combined with ganglion cyst excision in 21.43%. In our study, surgical treatment led to improvement in foot strength, with statistical relevance, in both sexes. Only one complication was observed.

Conclusions: Surgical exploration and repair of peroneal nerve injuries achieved good results in this series, with functional improvement of the analyzed patients in both sexes. When appropriate, surgical repair can lead to favorable outcome and early surgery can be a therapeutic strategy in selected cases.

Keywords: Peroneal nerve; Nerve graft; Micro neurolysis; Compression; Neuropathy

Introduction

Peroneal nerve injury is the most frequent pathology of the lower limbs, and accounts for about 15-20% of all peripheral nerve lesions [1]. The peroneal nerve arises from the roots of L4-S2 and is a lateral division of the sciatic nerve.

The sciatic nerve divides into the tibial and common peroneal nerves in the apex of the popliteal fossa. The common peroneal nerve subsequently travels around the fibular neck and then passes between the two heads of the peroneus longus muscle. The common peroneal nerve then divides into both the superficial and deep branches [2,3].

The location of the peroneal nerve, which is superficially located over the fibular head, is responsible for its vulnerability [1,4,5]. The peroneal nerve seems particularly prone to injury from motor vehicle accidents, gunshot wounds, iatrogenic accidents, and sport injuries [1]. Beyond that the compression of the peroneal nerve could be induced by a ganglion cyst, cysts of the lateral meniscus, or a tumor of the fibula head [2,6,7].

The peroneal nerve injury result in a variety of symptoms, including foot drop due to paralysis of the affected musculature, as well as sensory disturbances over the lateral surface of the lower extremity or the dorsum of the foot [2].

The diagnosis depends on understanding the anatomy of the peroneal nerve and on clinical findings. The literature recommends surgical decompression of the common peroneal nerve in some cases,

but surgical results are rarely described. Adequate surgical repair is responsible for the physiological maintenance of gait biomechanics [2].

Our objectives were to describe the neurosurgical technical nuances of peroneal nerve injury management and to analyze the outcomes of patients diagnosed with peroneal nerve injury and operated on by a single neurosurgeon at a private institution.

Materials and Methods

We performed a retrospective analysis with data collection from the medical records of patients diagnosed with peroneal nerve injury surgically treated by a single neurosurgeon at a single private institution, from 2000 to 2018. Thirty medical records were analyzed, and fourteen patients were included in the analysis. The inclusion criteria were patients >18 years old, peroneal nerve injury confirmed by electroneuromyography, patients whose medical records showed preoperative neurological status, and patients with a minimum follow-up of 6 months. Patients <18 years old, without electroneuromyographic studies, or patients with a follow-up of <6 months were not included in the study.

The variables analyzed were demographic characteristics, etiology of the lesion, preoperative neurological status, strength graduation (Table 1) [8], time between symptoms and surgery, location of lesion, perioperative finding, surgical technique (suture, neurolysis, and graft), complications, and neurological status 6 months after surgery.

Table 1: Muscle Grading System for Peroneal Nerve Injuries.

Grade	Indications
0	No palpable muscle contraction.
1	Palpable contraction of the peronei or anterior tibial muscles.
2	Peronei or anterior tibial muscles contract against gravity.
3	Peronei or anterior tibial muscles contract against gravity and give some resistance.
4	Peronei and anterior tibial muscles contract against moderate resistance.
5	Peronei and anterior tibial muscles contract against great resistance.

Kim et al. 2004 [8].

Table 2: Influence of sex on etiology.

Etiology	Sex		Total
	Female	Male	
Spontaneous	25% (1/4)	0% (0/10)	7.14% (1/14)
Traumatic	50% (2/4)	70% (7/10)	64.29% (9/14)
Tumoral	25% (1/4)	20% (2/10)	21.43% (3/14)
Mixed	0% (0/4)	10% (1/10)	7.14% (1/14)

The proportional data was presented with a descriptive analysis and all the non-parametric data was compared with the data from the Wilcoxon test; any value <0.05 was considered statistically significant. The utilized software was the Statistical Analyses System (SAS).

Results

Fourteen patients were analyzed, of which four were female and ten male. Table 2 demonstrates the influence of sex on etiology, especially in traumatic injuries, with 70% occurring in men and 50% in women. Tumoral lesions occurred in 25% of cases among female patients and in 20% among male patients. Mixed injuries were seen only in men with a prevalence of 10%.

Regardless of diagnose, 50% of the patients were diagnosed with traumatic neuroma (Figure 1) and 21.43% with ganglion cyst (Figure 2), as shown in Table 3.




Figure 1: A 64-year-old patient with traumatic neuroma, victim of car trauma with injury of the left common peroneal nerve. At admission, the patient presented with grade 0 strength on his left foot. He underwent neurolysis 3 months after trauma and evolved with complete strength recovery at 6-months follow-up.

Table 3: Influence of sex on preoperative diagnosis.

Diagnosis	Sex		Total
	Female	Male	
Fibrosis and Stenosis	0% (0/4)	10% (1/10)	7.14% (1/14)
Ganglion Cyst	0% (0/4)	30% (3/10)	21.43% (3/14)
Traumatic Neuroma	50% (2/4)	50% (5/10)	50% (7/14)
Schwannoma of Left Popliteal Fossa	25% (1/4)	0% (0/10)	7.14% (1/14)
Pale Looking Nerve	0% (0/4)	10% (1/10)	7.14% (1/14)
Fibrosis	25% (1/4)	0% (0/10)	7.14% (1/14)

Table 4: Influence of sex on the injury site.

Injury Site	Sex		Total
	Female	Male	
Left Common Peroneal	50% (2/4)	20% (2/10)	28.57% (4/14)
Right Common Peroneal	25% (1/4)	40% (4/10)	35.71% (5/14)
Right and Left Common Peroneal	25% (1/4)	0% (0/10)	7.14% (1/14)
Right Superficial Peroneal	0% (0/4)	10% (1/10)	7.14% (1/14)
Common Sciatic, Tibial, and Peroneal Rights	0% (0/4)	20% (2/10)	14.29% (2/14)
Left Common Sciatic and Peroneal	0% (0/4)	10% (1/10)	7.14% (1/14)

Table 5: Operative technique used according to sex.

Technique	Sex		Total
	Female	Male	
Graft	25% (1/4)	10% (1/10)	14.29% (2/14)
Neurolysis	50% (2/4)	50% (5/10)	50.00% (7/14)
Neurolysis and Graft	0% (0/4)	10% (1/10)	7.14% (1/14)
Exeresis	25% (1/4)	0% (0/10)	7.14% (1/14)
Ganglion Cyst Excision and Neurolysis	0% (0/4)	30% (3/10)	21.43% (3/14)

The most common injury observed in our series was isolated right common peroneal injury (35.71%), followed by isolated left common peroneal injury (28.75%), and then by joint nerve injury right sciatic, tibial and peroneal (14.29%), as shown in Table 4.

In our analysis, 42.85% of patients (6/14) underwent surgical treatment at less than 3 months after symptom onset (interval ranging from 7 days to 60 days), and 57.14% (8/14) underwent treatment 3 months after symptom onset (3 to 9 months). After analysis of patients who had complete strength recovery after 6 months of follow-up (42.85%; 6/14), 50% of these (3/6 patients) were approached within 3 months of symptom onset; all these patients showed blunt injuries and neurolysis was the technique performed, whether mixed or isolated.

The most commonly performed surgical technique was neurolysis, either isolated (50%), combined with a graft (7.14%), or combined with ganglion cyst excision (21.43%). Isolated graft (Figure 3) was performed in 14.29% of the cases and combined with other techniques in 7.14% of the cases (Table 5). In 3 cases in which the graft technique was used, all patients had grade 0 strength in the preoperative period and the results obtained at follow-up were little or no improvement in strength (33% Grade 0, 33% Grade 1, 33% Grade 3 at 6 months postoperatively).





Table 6: Influence of sex on time from symptom onset to surgery, preoperative strength and at 6 months after surgery.

Variable	Female			Male		
	Mean ± SD	Median	Variation	Mean ± SD	Median	Variation
Age (years)	30.0 ± 10.8	29.5	19 - 42	40.0 ± 15.11	40	12 - 64
Time from symptoms to surgery (days)	96.75 ± 96.28	100	7 - 180	91.50 ± 67.25 ^a	90	15 - 270
Preoperative strength	1.25 ± 2.50 ^{Aa}	0	0 - 5.0	0.80 ± 1.61 ^{Aa}	0	0.0 - 5.0
Strength 6 months after surgery	3.75 ± 1.50 ^{Ba}	4	2.0 - 5.0	3.00 ± 2.06 ^{Ba}	3	0.0 - 5.0

Means followed by different letters, lower case in the row and upper case in the column, differ (p <0.05; Wilcoxon test).



to surgery, preoperative strength, or strength 6 months after surgery (p >0.05). However, when strength was compared in preoperatively and at the 6-month follow-up, an increase was observed for both female and male patients (p <0.05) (Table 6).

Regarding complications of the procedures, only one male patient (7.14%; 1/14) presented surgical site infection in the postoperative period.

Discussion

In our study, traumatic injury was the most common cause of peroneal nerve injury, accounting for 64.27% of cases. This is similar to some results described in the literature [8-10], but contrasts with a study from the University of Louisiana, in which 87% of injuries were traumatic [8]. Traumatic injury could be explained by the susceptibility of the peroneal nerve to tensile forces due to its location and internal anatomy [8].

The main cause of tumoral lesions was ganglion cyst, corresponding to 21.43% of the cases, differing with that reported in the literature, of 40% [11]. The involvement of the peripheral nerves with ganglion cysts has been extensively studied and described in the literature, and multiple theories have been proposed, including cystic degeneration of neurilemmomas, expansion of synovial rests displaced during embryological development, herniation of joint synovium, and traumatic intraneural hemorrhage [12].

The lesion of the right common peroneal nerve was more common than that of the left (35.71%), as found in the study by Terzis et al., in which 64% of the injuries occurred also on the right common peroneal nerve (40 of 62 patients) [1].

Regarding treatment, isolated neurolysis was the most performed

The existence of pain in the preoperative period was described in 42.86% of the cases (6/14), 75% (3/4) among women and 30% (3/10) among men. Despite this, hyperesthesia or hypoesthesia was reported in 28.57% (4/14) and 35.71% (5/14) of cases, respectively. Among women, 50% (2/4) presented hyperesthesia, whereas among men, 20% (2/10) and 50% (5/10), presented hyperesthesia and hypoesthesia, respectively.

After a 6-month follow-up, pain was reported only in one woman (7.17%; 1/14); when sensitivity was assessed, 85.71% (12/14) were normal and hypoesthesia was described in two cases (14.29%; 2/14), including one patient of each sex.

Sex of the patient did not interfere in the time from symptom onset

procedure in our series, in 50% of the cases. This was similar to that reported in the study by Kim et al. [8], in which neurolysis was responsible for 38% of cases (121 patients out of a total of 318); and in the study by Terzis et al., in which 50% of the cases were treated with isolated micro neurolysis [1].

Repair, either by isolated graft or in conjunction with neurolysis, was performed in approximately 21% of our cases, compared with 43% [6] and 43.5% [1] in other studies. The graft sizes we used ranged from 5 to 8 cm and only one of three patients had partially recovered strength from grade 0 to grade 3, whereas the others did not benefit from the treatment. It has been well established that the functional recovery of patients with grafts is more difficult, especially those with more extensive grafts. Patients with smaller grafts generally have better outcomes, because there is a direct correlation between graft length and severity of injury [8].

In our study, surgical treatment allowed improvement in strength in both sexes, with statistical relevance ($p < 0.05$), as has also been shown in other centers [8,9]. Analysis of the patients with complete strength recovery after 6 months showed that 50% were approached within 3 months from symptom onset. The most accepted recommendation is that surgery should be performed as soon as possible on open injuries, but a delay of at least 3 months should be allowed on blunt injuries, due to the possibility of spontaneous recovery [1]. However, one of our patients, a victim of car accident, was approached within 2 months of injury, because he remained with significant loss of motor strength in the foot without any clinical or electroneuromyography improvement in that period of observation. He evolved in the postoperative period with complete strength recovery, showing that early surgery can be a therapeutic alternative in selected cases.

Conclusions

Our study has some limitations: it is a retrospective analysis, with a small number of patients treated in a single institution. We concluded that surgical exploration and repair of peroneal nerve injuries might improve patient outcomes. The goal of surgical treatment is to accomplish useful motor function. In selected cases, early treatment can be a therapeutic strategy.

References

1. Terzis JK, Kostas I. Outcomes with microsurgery of common peroneal nerve lesions. *J Plast Reconstr Aesthet Surg.* 2020; 73: 72-80.
2. Humphreys DB, Novak CB, Mackinnon SE. Patient outcome after common peroneal nerve decompression. *J Neurosurg.* 2007; 107: 314-318.
3. Wilson TJ, Maldonado AA, Amrami KK, Glazebrook KN, Moynagh MR, Spinner RJ. The anatomic location and importance of the tibialis posterior fascicular bundle at the sciatic nerve bifurcation: report of 3 cases. *J Neurosurg.* 2018; 131: 1869-1875.
4. Sunderland S. *Nerve and Nerve injuries.* 2nd ed. New York: Churchill Livingstone. 1978: 967-999.
5. Kretschmer T, Antoniadis G, Braun V, Rath SA, Richter HP. Evaluation of iatrogenic lesions in 722 surgically treated cases of peripheral nerve trauma. *J Neurosurg.* 2001; 94: 905-912.
6. Robin F, Kuchenbuch M, Sauleau P, Marleix S, Lucas G, Fraisse B, et al. [Peroneal nerve palsy in children: Uncommon diagnosis of a proximal tibiofibular synovial cyst]. *Arch Pediatr.* 2016; 23: 90-93.
7. Heilbrun ME, Tsuruda JS, Townsend JJ, Heilbrun MP. Intraneural perineurioma of the common peroneal nerve. Case report and review of the literature. *J Neurosurg.* 2001; 94: 811-815.
8. Kim DH, Murovic JA, Tiel RL, Kline DG. Management and outcomes in 318 operative common peroneal nerve lesions at the Louisiana State University Health Sciences Center. *Neurosurgery.* 2004; 54: 1421-1428; discussion 1428-1429.
9. Thoma A, Fawcett S, Ginty M, Veltri K. Decompression of the common peroneal nerve: experience with 20 consecutive cases. *Plast Reconstr Surg.* 2001; 107: 1183-1189.
10. Banerjee T, Koons DD. Superficial peroneal nerve entrapment. Report of two cases. *J Neurosurg.* 1981; 55: 991-992.
11. Nucci F, Artico M, Santoro A, Bardella L, Delfini R, Bosco S, et al. Intraneural synovial cyst of the peroneal nerve: report of two cases and review of the literature. *Neurosurgery.* 1990; 26: 339-344.
12. Harbaugh KS, Tiel RL, Kline DG. Ganglion cyst involvement of peripheral nerves. *J Neurosurg.* 1997; 87: 403-408.