

Special Article - Antibiotic Resistances

# Insights into the Topology of Antibody by Gamma Function

Liang L and Zhao B\*

College of Science, Northwest Agriculture and Forestry University, Yangling, Shaanxi, China

\*Corresponding author: Bin Zhao, College of Science, Northwest Agriculture and Forestry University, Yangling, China

Received: May 31, 2017; Accepted: July 14, 2017;

Published: July 21, 2017

**Abstract**

Along with the rapid development of genetic engineering technology and antibody engineering technology, humanized monoclonal antibody has been rapidly developed and gradually replaces the rat sourced monoclonal antibody. In this paper, we establish two new logarithmically completely monotonic functions involving the gamma function according to two preferred interaction geometries, necessary and sufficient conditions are presented for one of them to be logarithmically completely monotonic. As a consequence, a sharp inequality involving the gamma function is deduced to solve the problems of genetically engineered antibody.

**Keywords:** Gamma function; Genetically engineered antibody; Logarithmically completely monotonic; Inequality; Psi function

**Introduction**

Antibodies have been proven to be indispensable tools for biomedical applications. Different engineered antibodies have been developed for various purposes according to the amino acid sequence and/or spatial structure of protein (Figure 1). At present, it is still difficult to predict the optimal structure of antibodies. Topology knowledge can be important in antibody application as well as transformation. Theoretically, we can obtain desired antibodies by using protein/gene engineering technology. For instance, we can transform the Complementarity Determining Region (CDR) to promote the affinity of the antibody to antigen. Similarly, we could also transform any domain of antibody to make it bind with any desired target. Under this vision, topology is a powerful tool to predict the structure of protein and it will serve to antibody engineering. Our present work tries to explain, and predict, if possible, the change of structure, size and function of antibodies as well as their fragments from a topological perspective.

For  $\text{Re}(z) > 0$  the classical Euler's gamma function  $\Gamma$  and psi (digamma) function  $\Psi$  are defined by

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt, \psi(z) = \frac{\Gamma'(z)}{\Gamma(z)} \quad (1.1)$$

respectively. The derivatives  $\psi^{(n)}(z)$  for  $n \in \mathbb{N}$  are known as polygamma functions.


For  $\Psi(z)$  (see [1]), the following series representations are established:

$$\psi(z) = -\gamma - \frac{1}{z} + \sum_{n=1}^\infty \frac{z}{n(z+n)}, \quad (1.2)$$

$$\psi(z) = \ln z + \int_0^\infty \left( \frac{1}{t} - \frac{1}{1-e^{-t}} \right) e^{-zt} dt, \quad (1.3)$$

$$\psi(z) = -\gamma + \int_0^\infty \frac{e^{-t} - e^{-zt}}{1-e^{-t}} dt, \quad (1.4)$$

where  $\gamma = 0.577215664901$  denotes the Euler's constant. We next recall [2-5] that a function  $f$  is said to be completely monotonic on an interval  $I$ , if  $f$  has derivatives of all orders on  $I$  which alternate



**Figure 1:** Different antibody formats. a: different antibody or engineered antibodies; b: different shape of antibody.

successively in sign, that is,

$$(-1)^n f^{(n)}(x) \geq 0 \quad (1.5)$$

for all  $x \in I$  and for all  $n \geq 0$ . If inequality (1.5) is strict for all  $x \in I$  and all  $n \geq 0$ , then  $f$  is said to be strictly completely monotonic.



The classical Bernstein-Widder theorem [6, p. 160, Theorem 12a] states that a function  $f$  is completely monotonic on  $(0, \infty)$  if and only if it is a Laplace transform of some nonnegative measure  $\mu$ , that is,

$$f(x) = \int_0^\infty e^{-xt} d\mu(t), \quad (1.6)$$

where  $\mu(t)$  is non-decreasing and the integral converges for  $x > 0$ . We recall also [7-9] that a positive function  $f$  is said to be logarithmically completely monotonic on an interval  $I$  if  $f$  has derivatives of all orders on  $I$  and

$$(-1)^n [\ln f(x)]^{(n)} \geq 0, \quad (1.7)$$

for all  $x \in I$  and for all  $n \geq 1$ . If inequality (1.7) is strict for all  $x \in I$  and all  $n \geq 1$ , then  $f$  is said to be strictly logarithmically completely monotonic. The antibody structure will be changed when it binds certain target (Figure 2a), i.e.: antigen, receptor. How to describe



the changes in the view of topology? The following cases will explain it in detail. It was proved explicitly in [8] and other articles that a logarithmically completely monotonic function must be completely monotonic. In [10], G. D. Anderson et al. proved that the function

$$g(x) = x(\ln x - \psi(x)) \tag{1.8}$$

is strictly decreasing and strictly convex on  $(0, \infty)$ , with two limits

$$\lim_{x \rightarrow \infty} g(x) = 1, \lim_{x \rightarrow \infty} g'(x) = \frac{1}{2} \tag{1.9}$$

From (1.9) and the monotonicity of  $g(x)$ , then the double inequalities

$$\frac{1}{2x} < \ln x - \psi(x) < \frac{1}{x} \tag{1.10}$$

holds for all  $x > 0$ .

In [11, Theorem 1], by using the well-known Binet's formula, H. Alzer generalized the monotonicity and convexity of  $g(x)$ , that is, the function

$$g_\alpha(x) = x^\alpha (\ln x - \psi(x)) \tag{1.11}$$

is strictly completely monotonic on  $(0, \infty)$  if and only if  $\alpha \leq 1$ .

In [12], D. Kershaw and A. Laforgia proved that the function  $[\Gamma(1+1/x)]^x$  is decreasing on  $(0, \infty)$  and  $x[\Gamma(1+1/x)]^x$  is increasing on  $(0, \infty)$ . These are equivalent to the function  $[\Gamma(1+1/x)]^x$  being increasing and  $[\Gamma(1+1/x)]^x / x$  being decreasing on  $(0, \infty)$ , respectively.

In [13], Theorem 5], F. Qi and Ch.-p. Chen generalized these functions. They obtained the fact that for all  $x > 0$  the function  $x^r [\Gamma(1+x)]^{1/x}$  is strictly increasing for  $r \geq 0$  and strictly decreasing for  $r \leq -1$ , respectively.

After the papain digestion, the remained antibody functional part (usually the Fab domain), will be smaller and the structure is also changed (Figure 1b). These changes can be revealed vividly using

topology. Recently [[14], Theorem 1], F. Qi, C.-F Wei and B.-N Guo established another excellent result, which states that for given  $y \in (-1, \infty)$  and  $\alpha \in (-\infty, \infty)$  let

$$g_{\alpha,y}(x) = \begin{cases} \frac{1}{(x+y+1)^\alpha} \left[ \frac{\Gamma(x+y+1)}{\Gamma(y+1)} \right]^{1/x}, & x \in (-y-1, \infty) \setminus \{0\}; \\ \frac{1}{(y+1)^\alpha} \exp[\Psi(y+1)], & x = 0. \end{cases} \tag{1.12}$$

The function (1.12) is logarithmically completely monotonic with respect to  $x \in (-y-1, \infty)$  if and only if  $\alpha \geq \max\{1, 1/(y+1)\}$ ; and if  $\alpha \leq \min\{1, 1/2(y+1)\}$ ; the reciprocal of the function (1.12) is logarithmically completely monotonic with respect to  $x \in (-y-1, \infty)$ .

Antibodies occur spontaneously gathering and forming dimer, polymer, which will influence their functions (Figure 2b). In antibody engineering practice, it urgently needs some measures to overcome this difficulty. From topology perspective, we could understand this issue as follow.

Stimulated by the above results, we put forward the function as follows: for given  $y \in (0, \infty)$  and real number  $\alpha$ , let the function  $f_{\alpha,y}(x)$  defined by

$$f_{\alpha,y}(x) = \frac{1}{x^\alpha} \left[ \frac{\Gamma(x+y)}{\Gamma(y)} \right]^{1/x}, \quad x \in (-y, \infty) \setminus \{0\}. \tag{1.13}$$

Our first result is contained in the following theorem.

**Theorem 1:** For the function (1.13), then the following statements are true:

- (1) For any given  $y \geq 1$ , the function (1.13) is strictly logarithmically completely monotonic with respect to  $x \in (-y, \infty) \setminus \{0\}$  if and only if  $\alpha \geq 1$ ;
- (2) For any given  $0 < y < 1$ , if  $\alpha \geq e^{(1-y)}/y$  then the function (1.13) is strictly Logarithmically completely monotonic with respect to  $x \in (-y, \infty) \setminus \{0\}$ ;
- (3) For any given  $y > 0$ , the reciprocal of the function (1.13) is strictly

Logarithmically completely monotonic with respect to  $x \in (-y, \infty) \setminus \{0\}$  if and only if  $\alpha \geq 0$ .

Our second result is presented in the following theorem.

**Theorem 2:** For any given  $y \in [1, \infty)$ , let the function  $h_y(x)$  be defined on  $(0, \infty)$  by

$$h_y(x) = \frac{(x+y)^x}{\Gamma(x+y)x^{x^2}} e^{\left(\frac{1}{2}-\gamma\right)x^2-2x} \exp\left(\sum_{n=1}^{\infty} \left(\frac{x^2}{n} + \ln\left(1 + \frac{x}{n}\right) - 2x\right)\right) \tag{1.14}$$

where  $\gamma$  denotes the Euler's constant, then the function (1.14) is strictly logarithmically completely monotonic with respect to  $x$  on  $(0, \infty)$ .

The following corollary can be derived from Theorems 2 immediately.

**Corollary 1:** For any given  $y \geq 1$ , the inequality

$$\frac{(x+y)^x}{\Gamma(x+y)x^{x^2}} e^{\left(\frac{1}{2}-\gamma\right)x^2-2x} < \frac{1}{\Gamma(y)} \exp\left(\sum_{n=1}^{\infty} \left(\frac{x^2}{n} + \ln\left(1 + \frac{x}{n}\right) - 2x\right)\right) \tag{1.15}$$

holds for all  $x > 0$ .

**Lemma**

In order to prove our main results, we need the following lemmas. It is well known that Bernoulli polynomials  $B_k(x)$  and Euler polynomials  $E_k(x)$  are defined by

$$\frac{te^{tx}}{e^t-1} = \sum_{n=0}^{\infty} \frac{B_n(x)}{n!} t^n, |t| < 2\pi, \tag{2.1}$$

$$\frac{2e^{tx}}{e^t+1} = \sum_{n=0}^{\infty} \frac{E_n(x)}{n!} t^n, |t| < \pi, \tag{2.2}$$

respectively. The Bernoulli numbers  $B_n$  are denoted by  $B_n = B_n(0)$ , while the Euler numbers  $E_n$  are defined by  $2^n E_n(1/2)$ .

In [15,16], the following summation formula is given:

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^{2k+1}} = \frac{\pi^{2k+1} E_k}{2^{2k+1} (2k)!} \tag{2.3}$$

for any nonnegative integer  $k$ , which implies

$$B_{2n} = (-1)^{n-1} \frac{2(2n)!}{(2\pi)^{2n}} \sum_{m=1}^{\infty} \frac{1}{m^{2n}}, n \in \mathbb{N}. \tag{2.4}$$

In particular, it is known that for all  $n \in \mathbb{N}$  we have

$$B_{2n+1} = 0, B_{2n} = (-1)^{n+1} |B_{2n}|, \tag{2.5}$$

$$E_{2n+1} = 0, E_{2n} = (-1)^n |E_{2n}|, \tag{2.6}$$

And the first few nonzero values are

$$B_0=0, B_1 = -\frac{1}{2}, B_2 = \frac{1}{6}, B_4 = -\frac{1}{30},$$

$$E_0=1, E_2=-1, E_4=5$$

(See [[17], p.804, Chapter23]).

The Bernoulli and Euler numbers and polynomials are generalized (see [18-21]).

**Lemma 1:** (see [22,23]) For real number  $x > 0$  and natural number, then

$$\ln \Gamma(x) = \ln \sqrt{2\pi} + \left(x - \frac{1}{2}\right) \ln x - x + \sum_{n=1}^m \frac{B_{2n}}{2(2n-1)n} \frac{1}{x^{2n-1}} + \tag{2.7}$$

$$\theta_1 \frac{B_{2(m+1)}}{(2m+1)(2m+2)} \cdot \frac{1}{x^{2m+1}}, 0 < \theta_1 < 1;$$

$$\Psi(x) = \ln x - \frac{1}{2x} - \sum_{n=1}^m \frac{B_{2n}}{2n} \cdot \frac{1}{x^{2n}} - \theta_2 \cdot \frac{B_{2(m+1)}}{2m+2} \cdot \frac{1}{x^{2m+2}}, 0 < \theta_2 < 1; \tag{2.8}$$

$$\Psi'(x) = \frac{1}{x} + \frac{1}{2x^2} + \sum_{n=1}^m \frac{B_{2n}}{x^{2n+1}} + \theta_3 \cdot \frac{B_{2(m+1)}}{x^{2m+3}}, 0 < \theta_3 < 1;$$

$$\Psi''(x) = -\frac{1}{x^2} - \frac{1}{x^3} - \tag{2.9}$$

$$\sum_{n=1}^m \frac{B_{2n}}{x^{2n+1}} - (2m+3)\theta_4 \cdot \frac{B_{2(m+1)}}{x^{2m+4}}, 0 < \theta_4 < 1. \tag{2.10}$$

**Remark 1:**  $\theta_1, \theta_2, \theta_3, \theta_4$  only depend on natural number  $m$ .

**Lemma 2:** (see [[24-31], p. 107, Lemma 3]) For real number  $x > 0$  and natural number  $n$ , we have

$$\frac{(n-1)!}{x^n} + \frac{n!}{2x^{n+1}} < (-1)^{n+1} \Psi^{(n)}(x) < \frac{(n-1)!}{x^n} + \frac{n!}{x^{n+1}}. \tag{2.11}$$

**Lemma 3:** (see [1,17]) For real number  $x > 0$  and natural number  $n$ , we have

$$\ln x = \int_0^{\infty} \frac{e^{-t} - e^{-xt}}{t} dt, \tag{2.12}$$

$$\frac{\Gamma(n)}{x^n} = \int_0^{\infty} t^{n-1} e^{-xt} dt, \tag{2.13}$$

$$\Psi^{(n)}(x) = (-1)^{n+1} \int_0^{\infty} \frac{t^n e^{-xt}}{1 - e^{-t}} dt. \tag{2.14}$$

**Lemma 4:** Let the sequence of functions  $u_n(x)$  for  $n \in \mathbb{N}$  be defined on  $[0, \infty)$  by

$$u_n(x) = \frac{x^2}{n} + \ln\left(1 + \frac{x}{n}\right) - 2x, \tag{2.15}$$

the series  $\sum_{n=1}^{\infty} u_n(x)$  is differentiable on  $[0, \infty)$ , that is,

$$\left(\sum_{n=1}^{\infty} u_n(x)\right)' = \sum_{n=1}^{\infty} \left(\frac{2x}{n} - \frac{2x}{n+x}\right). \tag{2.16}$$

**Proof:** It is obvious that  $\sum_{n=1}^{\infty} u_n(0) = 0$ , therefore  $\sum_{n=1}^{\infty} u_n(x)$  converges at  $x=0$ . In order to prove (2.16), we need only to show that the inner closed uniform convergence of the  $\sum_{n=1}^{\infty} u_n'(x)$  on  $[0, \infty)$ . From (2.15), we have

$$\sum_{n=1}^{\infty} u_n'(x) = \sum_{n=1}^{\infty} \left(\frac{2x}{n} - \frac{2x}{n+x}\right). \tag{2.17}$$

For any interval  $[a,b] \subset [0, \infty)$ , we have

$$\left|\sum_{n=1}^{\infty} \left(\frac{2x}{n} - \frac{2x}{n+x}\right)\right| \leq \sum_{n=1}^{\infty} \frac{2b^2}{n^2} \tag{2.18}$$

For all  $x \in [a,b]$ . It is easy to check that the series  $\sum_{n=1}^{\infty} \frac{2b^2}{n^2}$  converges, which and Weierstrass M-test implies that the series  $\sum_{n=1}^{\infty} u_n'(x)$  is inner closed uniformly convergent on  $[0, \infty)$ . Hence the series  $\sum_{n=1}^{\infty} u_n(x)$  is differentiable on  $[0, \infty)$  and the identity (2.17) holds for  $x \geq 0$ .

The lemma is proved.

**Lemma 5:** For  $0 < a \leq 1$  and real number  $b$ , let the function  $Q_{a,b}(x)$  be defined by

$$Q_{a,b}(x) = \frac{(ax+b)}{\Gamma(ax+b)}, x \in \left(-\frac{b}{a}, \infty\right) \tag{2.19}$$

If  $b \geq a$ , then the function (2.19) satisfies

$$(-1)^n [\ln Q_{a,b}(x)]^{(n)} > 0 \tag{2.20}$$

for all  $x \in \left(-\frac{b}{a}, \infty\right)$  and  $n = 2, 3, \dots$ .

**Proof:** Taking the logarithm of  $Q_{a,b}(x)$  yields

$$\ln Q_{a,b}(x) = x \ln(ax+b) - \ln \Gamma(ax+b), \tag{2.21}$$

and differentiating  $\ln Q_{a,b}(x)$ , then

$$(\ln Q_{a,b}(x))' = x \ln(ax+b) - a\Psi(ax+b) - \frac{b}{ax+b} + 1. \tag{2.22}$$

For given integer  $n \geq 2$ , we get

$$(\ln Q_{a,b}(x))^{(n)} = \frac{(-1)^n (n-2)! a^{n-1}}{(ax+b)^{n-1}} \tag{2.23}$$

and, by the identities (2.13) and (2.14), (2.23) can be written as

$$a^n \Psi^{(n-1)}(ax+b) - \frac{(-1)^n (n-1)! b a^{n-1}}{(ax+b)^n} \tag{2.24}$$

$$(\ln Q_{a,b}(x))^{(n)} = (-1)^n a^{n-1} \int_0^\infty \frac{e^{-(ax+bt)} t^{n-2}}{1-e^{-t}} \left( (1+bt)(1-e^{-t}) - at \right) dt.$$

Let  $p(t) = (1+bt)(1-e^{-t}) - at$

and  $q(t) = (1+t)(1-e^{-t}) - t$

It is easy to check that

$$q'(t) = te^{-t} > 0, t \in (0, \infty) \tag{2.25}$$

Therefore  $q(t)$  is strictly increasing  $[0, \infty)$ , and then  $q(t) > q(0) = 0$ . The following two cases will complete the proof of Lemma 5.

**Case 1:** If  $0 < a \leq 1 \leq b$ , then since  $q(t) > 0$  for  $t > 0$ , we have

$$at \leq (1+bt)(1-e^{-t}) \leq (1+bt)(1-e^{-t}), \tag{2.26}$$

which implies  $at < (1+bt)(1-e^{-t})$ , and then  $P(t) > 0$  for all  $t > 0$ . From (2.24), we know that the inequality (2.20) holds for  $x \in (-b/a, \infty)$  and integer  $n \geq 2$ .

**Case 2:** If  $0 < a \leq b \leq 1$  then we get

$$p'(t) = b - a + e^{-t}(bt + 1 - b) \geq \tag{2.27}$$

$$bte^{-t} > 0, t \in (0, \infty),$$

Therefore  $P(t)$  is strictly increasing on  $(0, \infty)$ , and then  $P(t) > P(0) = 0$ .

From (2.24), we know that the inequality (2.20) holds for  $x \in (-b/a, \infty)$  and integer  $n \geq 2$ .

The lemma is proved.

### Proof of Theorems

#### Proof of Theorem 1

For  $x \neq 0$  and natural number  $n$ , taking the logarithmically

differential into consideration yields

$$[f_{\alpha,y}(x)]^{(n)} = \frac{1}{x^{n+1}} \left[ \sum_{k=0}^n (-1)^k \frac{n! x^{n-k} \Psi^{(n-k)}(x+y)}{(n-k)!} - (-1)^n n! \ln \Gamma(y) - \alpha (-1)^{n-1} (n-1)! x \right] \tag{3.1}$$

Where  $\Psi^{(-1)}(x+y)$  and  $\Psi^{(0)}(x+y)$  stand for  $\ln \Gamma(x+y)$  and  $\psi(x+y)$  respectively.

Furthermore, differentiating  $x^{n+1} [f_{\alpha,y}(x)]^{(n)}$  directly gives

$$\left\{ x^{n+1} [f_{\alpha,y}(x)]^{(n)} \right\}' = (-1)^{n-1} x^n \left[ (-1)^{n+1} \Psi^{(n)}(x+y) - \frac{\alpha(n-1)!}{x^n} \right]. \tag{3.2}$$

Making use of (2.11) and (2.13) shows that for all  $n \in \mathbb{N}$  and any fixed  $y > 0$ , the double inequality

$$\int_0^\infty t^{n-1} e^{-xt} \left[ e^{-yt} \left( 1 + \frac{1}{2}t \right) - \alpha \right] dt = \tag{3.3}$$

$$\frac{(n-1)!}{(x+y)^n} + \frac{1}{2} \frac{n!}{(x+y)^{n+1}} - \frac{\alpha(n-1)!}{x^n} \leq$$

$$\frac{(-1)^{n-1}}{x^n} \left\{ x^{n+1} [f_{\alpha,y}(x)]^{(n)} \right\} \leq \frac{(n-1)!}{(x+y)^n} + \frac{n!}{(x+y)^{n+1}} -$$

$$\frac{\alpha(n-1)!}{x^n} = \int_0^\infty t^{n-1} e^{-xt} [e^{-yt}(1+t) - \alpha] dt$$

holds for all  $x \in (-y, \infty) \setminus \{0\}$  and  $\alpha \in (-\infty, \infty)$  For any fixed  $y \in (0, \infty)$  let  $u(t)$  and  $v(t)$  be defined on  $(-\infty, \infty)$  by

$$u(t) = e^{-yt} \left( 1 + \frac{1}{2}t \right) \text{ and } v(t) = e^{-yt} (1+t) \text{ respectively.}$$

Differentiating  $u(t)$  and  $v(t)$  directly, we obtain

$$u'(t) = e^{-yt} \left( \frac{1}{2} - y - \frac{1}{2}yt \right), \tag{3.4}$$

$$v'(t) = e^{-yt} (1 - y - yt). \tag{3.5}$$

Therefore, for given  $y \in (0, \infty)$  we have

$$u'(t) \begin{cases} > 0, t < 1/y - 2, \\ < 0, t > 1/y - 2 \end{cases} \tag{3.6}$$

and

$$v'(t) \begin{cases} > 0, t < 1/y - 1, \\ < 0, t > 1/y - 2 \end{cases} \tag{3.7}$$

From (3.6) and (3.7), we conclude that for all  $t > 0$  we obtain.

$$u(t) > 0 \tag{3.8}$$

and

$$v(t) \begin{cases} < 1, y \geq 1, \\ \leq e^{-(1-y)/y}, 0 < y < 1 \end{cases} \tag{3.9}$$

From (3.3) and (3.8)-(3.9), it is easy to see that

$$\frac{(-1)^{n-1}}{x^n} \left\{ x^{n+1} [f_{\alpha,y}(x)]^{(n)} \right\}' \begin{cases} > 0, \text{ if } \alpha \leq 0 \text{ for fixed } y > 0 \\ < 0, \text{ if } \alpha \geq 1 \text{ for fixed } y \geq 1 \\ < 0, \text{ if } \alpha \geq e^{-(1-y)/y} \text{ for fixed } 0 < y < 1 \end{cases} \tag{3.10}$$

for all  $n \in \mathbb{N}$  and all  $x \in (-y, \infty) \setminus \{0\}$ .

On the one hand, if  $x \in (0, \infty)$  then the inequalities (3.10) can be equivalently changed into

$$\left\{ x^{2k+1} [f_{\alpha,y}(x)]^{(2k)} \right\}' \begin{cases} < 0, \text{ if } \alpha \leq 0 \text{ for fixed } y > 0 \\ > 0, \text{ if } \alpha \geq 1 \text{ for fixed } y \geq 1 \\ < 0, \text{ if } \alpha \geq e^{-(1-y)/y} \text{ for fixed } 0 < y < 1 \end{cases} \tag{3.11}$$

and

$$\left\{ x^{2k} [f_{\alpha,y}(x)]^{(2k-1)} \right\}' \tag{3.12}$$

$$\begin{cases} >0, \text{ if } \alpha \leq 0 \text{ for fixed } y > 0 \\ <0, \text{ if } \alpha \geq 1 \text{ for fixed } y \geq 1 \\ >0, \text{ if } \alpha \geq e^{-(1-y)} / y \text{ for fixed } 0 < y < 1 \end{cases}$$

For  $k \in \mathbb{N}$ .

From (3.1), then simple computation shows that

$$\lim_{x \rightarrow 0} x^{n+1} [f_{\alpha,y}(x)]^{(n)} = 0 \tag{3.13}$$

for all  $n \in \mathbb{N}$  and any given  $y \in (0, \infty)$ . As a result,

$$\begin{cases} x^{2k+1} [f_{\alpha,y}(x)]^{(2k)} \\ <0, \text{ if } \alpha \leq 0 \text{ for fixed } y > 0 \\ >0, \text{ if } \alpha \geq 1 \text{ for fixed } y \geq 1 \\ >0, \text{ if } \alpha \geq e^{-(1-y)} / y \text{ for fixed } 0 < y < 1 \end{cases} \tag{3.14}$$

and

$$\begin{cases} x^{2k} [f_{\alpha,y}(x)]^{(2k-1)} \\ >0, \text{ if } \alpha \leq 0 \text{ for fixed } y > 0 \\ <0, \text{ if } \alpha \geq 1 \text{ for fixed } y \geq 1 \\ <0, \text{ if } \alpha \geq e^{-(1-y)} / y \text{ for fixed } 0 < y < 1 \end{cases} \tag{3.15}$$

for all  $k \in \mathbb{N}$  and all  $x > 0$ .

Therefore, (3.14) and (3.15) imply

$$\begin{cases} (-1)^n [f_{\alpha,y}(x)]^{(n)} \\ <0, \text{ if } \alpha \leq 0 \text{ for fixed } y > 0 \\ >0, \text{ if } \alpha \geq 1 \text{ for fixed } y \geq 1 \\ >0, \text{ if } \alpha \geq e^{-(1-y)} / y \text{ for fixed } 0 < y < 1 \end{cases} \tag{3.16}$$

for all  $n \in \mathbb{N}$  and all  $x > 0$ . Hence, if either  $\alpha \geq e^{-(1-y)} / y$  for given  $0 < y < 1$  or  $\alpha \geq 1$  for given  $y \geq 1$ , the function (1.13) is strictly logarithmically completely monotonic with respect to  $x$  on  $(0, \infty)$ , and if  $\alpha \leq 0$  for given  $y > 0$ , so is the reciprocal of the function (1.13). On the other hand, if  $x \in (-y, 0)$  for any given  $y > 0$ , then (3.10) implies

$$\begin{cases} \left\{ x^{n+1} [f_{\alpha,y}(x)]^{(n)} \right\}' \\ <0, \text{ if } \alpha \leq 0 \text{ for fixed } y > 0 \\ >0, \text{ if } \alpha \geq 1 \text{ for fixed } y \geq 1 \\ >0, \text{ if } \alpha \geq e^{-(1-y)} / y \text{ for fixed } 0 < y < 1 \end{cases} \tag{3.17}$$

for  $n \in \mathbb{N}$ .

In view of (3.13), we can conclude that

$$\begin{cases} x^{n+1} [f_{\alpha,y}(x)]^{(n)} \\ >0, \text{ if } \alpha \leq 0 \text{ for fixed } y > 0 \\ <0, \text{ if } \alpha \geq 1 \text{ for fixed } y \geq 1 \\ <0, \text{ if } \alpha \geq e^{-(1-y)} / y \text{ for fixed } 0 < y < 1 \end{cases} \tag{3.18}$$

For  $n \in \mathbb{N}$ . It is obvious that (3.18) is equivalent to that (3.14) and (3.15) hold for any given  $y > 0$  and  $x \in (-y, 0)$ . Therefore, it is easy to prove similarly that (3.16) is also valid on  $x \in (-y, 0)$  for any given  $y > 0$  and all  $n \in \mathbb{N}$ . The amino acid of antibody/protein possesses different preferences. Thus we can conduct site-directed mutation to promote the affinity and/or hydrophilic with the prediction of topology. For example, bovine antibodies have an unusual structure comprising a  $\beta$ -strand 'stalk' domain and a disulphide-bonded 'knob' domain in CDR3 (Figure 2). Attempts have been made to utilize such amino acid preference for antibody drug development.

Consequently, the function (1.13) is the same logarithmically completely monotonicity on  $(-y, 0)$  as on  $(0, \infty)$ , that is, if either  $\alpha \geq e^{-(1-y)}$

$/y$  for given  $0 < y < 1$  or  $\alpha \geq 1$  for given  $y \geq 1$ , the function (1.13) is strictly logarithmically completely monotonic with respect to  $x$  on  $(-y, 0)$ , and if  $\alpha \leq 0$  for given  $y > 0$ , so is the reciprocal of the function (1.13).

Conversely, we assume that the reciprocal of the function (1.13) is strictly logarithmically completely monotonic on  $(-y, \infty) \setminus \{0\}$  for any given  $y > 0$ . There we have for any given  $y > 0$  and all  $x > 0$ .

$$f'_{\alpha,y}(x) = \frac{\ln \Gamma(x+y) - x\Psi(x+y) - \ln \Gamma(y) + \frac{\alpha}{x}}{x^2} < 0, \tag{3.19}$$

Which implies

$$\alpha < -\frac{\ln \Gamma(x+y) - x\Psi(x+y) - \ln \Gamma(y)}{x}. \tag{3.20}$$

By L'Hospital's rule, we have

$$\lim_{x \rightarrow 0} \frac{-\ln \Gamma(x+y) + x\Psi(x+y) + \ln \Gamma(y)}{x} = 0 \tag{3.21}$$

for any given  $y > 0$ . By virtue of (3.20) and (3.21), we conclude that the necessary condition for the reciprocal of the function (1.13) to be strictly logarithmically completely monotonic is  $\alpha \leq 0$ .

If the function (1.13) is logarithmically completely monotonic  $(-y, \infty) \setminus \{0\}$  for any given  $y > 0$ , then the inequality (3.19) and (3.20) are reversed for any given  $y > 0$  and all  $x > 0$ .

By utilizing (2.7) and (2.8), it is easy to see that

$$\lim_{x \rightarrow \infty} \frac{-\ln \Gamma(x+y) + x\Psi(x+y) + \ln \Gamma(y)}{x} = 1 \tag{3.22}$$

for any given  $y > 0$ . In fact, it is not difficult to show that the necessary condition for the function (1.13) to be strictly logarithmically completely monotonic is  $\alpha \geq 1$ .

The proof of theorem 1 is completed.

### Proof of Theorem 2

Taking the logarithm of  $h_y(x)$  gives

$$\begin{aligned} \ln h_y(x) &= \ln \frac{(x+y)^n}{\Gamma(x+y)} - \ln x^{x^2} + \\ &\left(\frac{1}{2} - \gamma\right)x^2 - 2x + \sum_{n=1}^{\infty} \left(\frac{x^2}{n} + \ln\left(1 + \frac{x}{n}\right) - 2x\right). \end{aligned} \tag{3.23}$$

$$\mu(x) = \ln \frac{(x+y)^x}{\Gamma(x+y)}, \tag{3.24}$$

$$\omega(x) = -\ln x^{x^2} + \left(\frac{1}{2} - \gamma\right)x^2 - \tag{3.25}$$

$$2x + \sum_{n=1}^{\infty} \left(\frac{x^2}{n} + \left(1 + \frac{x}{n}\right)^{2n} - 2x\right)$$

then

$$\ln h_y(x) = \mu(x) + \omega(x). \tag{3.26}$$

In view of Lemma 4, straightforward calculation gives

$$(\ln h_y(x))' = \ln(x+y) + \tag{3.27}$$

$$\frac{x}{x+y} + -\Psi(x+y) - 2t \ln t + 2\gamma x +$$

$$2 - \sum_{n=1}^{\infty} \left(\frac{2x}{n} - \frac{2x}{n+x}\right) = \mu'(x) + \omega'(x).$$

By virtue of (1.12), the identity (3.27) is equivalent to

$$(\ln h_y(x))' = \ln(x+y) + \tag{3.28}$$

$$\frac{x}{x+y} + -\Psi(x+y) -$$

$$2x(\ln x - \Psi(x)) = \mu'(x) + \omega'(x).$$

By Lemma 5, we know that  $\mu'(x)$  is strictly increasing on  $(0, \infty)$ ,



which and (1.10) imply the limit of  $\mu'(x)$  equals 1 as  $x \rightarrow \infty$ , therefore

$$\mu'(x) < 1 \quad (3.29)$$

holds for all. We know that  $g(x)$  is strictly completely monotonic on  $(0, \infty)$ , where  $g(x)$  is defined by (1.8), hence for given integer  $n \geq 0$ , the inequality

$$(-1)^{n+1} (\omega'(x))^{(n)} > 0 \quad (3.30)$$

holds for all  $x > 0$ .

And then by using inequality (1.9) and (1.10), we get

$$-2 < \omega'(x) < -1 \quad (3.31)$$

For all  $x > 0$ . From (3.29) and (3.31), we conclude that

$$(\ln h_y(x))' = \mu'(x) + \omega'(x) < 0 \quad (3.32)$$

for all  $x > 0$ . Utilizing Lemma 5 and (3.30), for given Integer  $n \geq 2$ , it is easy to see that

$$(-1)^n (\ln h_y(x))^{(n)} = (-1)^n (\mu(x))^{(n)} + (-1)^n (\omega(x))^{(n)} > 0 \quad (3.33)$$

for all  $x > 0$ .

Theorem 2 follows from (3.32) and (3.33).

Thus the Proof of Theorem 2 is completed.

## Conclusion

In conclusion we establish two new logarithmically completely monotonic functions involving the gamma function according to two preferred interaction geometries, and a sharp inequality involving the gamma function is deduced to solve the problems of genetically engineering antibody. It is necessary to address, many other aspects (such as thermal condition, alkalinity or acidity, adhesion of antibody) are also playing key roles in antibody functioning, which could be also understood from bio-mathematical perspective, and such knowledge will be in return useful for biomedical application of antibodies as well as proteins.

## Acknowledgements

This work was supported by the Fundamental Research Funds for the Central Universities (2014YB030), Ministry of Education and State Administration of Foreign Experts Affairs "Overseas Teacher" project (MS2011XBNL057), the Key Construction Program (2015SD0018) of International Cooperation Base in S&T, Shaanxi Province, China.

## References

- Erdélyi A, Magnus W, Oberhettinger F, Tricomi FG. Higher Transcendental Functions. Vol. I, II, III, McGraw-Hill. New York. 1953-1955.
- Bernstein S. Sur les fonctions absolument monotones. Acta Math. 1929; 52: 1-66.
- Hausdorff F. Summationsmethoden und Momentfolgen I. Math. Z. 1921; 9: 74-109.
- Qi F, Chen Ch-p. A complete monotonicity property of the gamma function, J. Math. Anal. Appl. 2004; 296: 603-607.
- Widder DV. Necessary and sufficient conditions for the representation of a function as a Laplace integral. Trans. Amer. Math. Soc. 1931; 33: 851-892.
- Widder DV. The Laplace Transform, Princeton University Press, Princeton. 1941.
- Atanassov RD, Tsoukrovski UV. Some properties of a class of logarithmically completely monotonic functions. C. R. Acad. Bulgare Sci. 1988; 41: 21-23.
- Berg C. Integral representation of some functions related to the gamma function. Mediterr. J. Math. 2004; 1: 433-439.
- Grinshpan AZ, Ismail MEH. Completely monotonic functions involving the gamma and q-gamma functions, Proc. Amer. Math. Soc. 2006; 134: 1153-1160.
- Anderson GD, Barnard RW, Richards KC, Vamanamurthy MK, Vuorinen M. Inequalities for zero-balanced hypergeometric functions. Trans. Amer. Math. Soc. 1995; 347: 1713-1723.
- Alzer H. On some inequalities for the gamma and psi functions. Math. Comp. 1997; 66: 373-389.
- Kershaw D, Laforgia A. Monotonicity results for the gamma function. Atti Accad. Sci. To-rino Cl. Sci. Fis. Mat. Natur. 1985; 119: 127-133.
- Qi F, Chen Ch-p. Monotonicity and convexity results for functions involving the gamma function. J. Appl. Math. Sci. 2004; 1: 27-36.
- Qi F, Wei C-F, Guo B-N. Complete monotonicity of a function involving the ratio of gamma functions and applications. Banach J. Math. Anal. 2012; 6: 35-44.
- Guo B-N, Qi F. A property of logarithmically absolutely monotonic functions and the logarithmically complete monotonicity of a power-exponential function. Politehn. Univ. Bucharest Sci. Ser. A Appl. Math. Phys. 2010; 72: 21-30.
- Wang Z-X, Guo D-R. Introduction to Special Function. The Series of Advanced Physics of Peking University. Peking University Press. Beijing. China. 2000.
- Abramowitz M, Stegun IA. Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables. 9th edition, Applied Mathematics Series, vol.55, National Bureau of Standards, Washington. 1970.
- Guo B-N, Qi F. "Generalization of Bernoulli polynomials". International Journal of Mathematical Education in Science and Technology. 2002; 33: 428-431.
- Luo Q-M, Guo B-N, Qi F, Debnath L. Generalizations of Bernoulli numbers and polynomials. International Journal of Mathematics and Mathematical Sciences, 2003; 2003: 3769-3776.
- Luo Q-M, Qi F. "Relationships between generalized Bernoulli numbers and polynomials and generalized Euler numbers and polynomials". Advanced Studies in Contemporary Mathematics (Kyungshang). 2003; 7: 11-18.
- Luo Q-M, Qi F, Debnath L. Generalizations of Euler numbers and polynomials. International Journal of Mathematics and Mathematical Sciences. 2003; 2003: 3893-3901.
- Qi F, Guo B-N. Monotonicity and convexity of ratio between gamma functions to different powers. Journal of the Indonesian Mathematical Society. 2005; 11: 39-49.
- Fichtenholz GM. Differential- und Integralrechnung, Vol. II, Berlin. VEB Deutscher Verlag der Wissenschaften. 1964.
- Guo B-N, Qi F. Two new proofs of the complete monotonicity of a function involving the psi function. Bull. Korean Math. Soc. 2010; 47: 103-111.
- Scott AM, Wolchok JD, Lloyd J. Antibody therapy of cancer [ J ]. Nature reviews. 2012; 12: 278- 287.
- Carlos F, Lucia R, Jeffrey W. High-affinity self-reactive human antibodies by design and selection: Targeting the integrin ligand binding site. Medical Sciences. 1993; 90:10003-10007.
- Miroslava O, Xavier E. Human genetics moves from clinic to bench - and back [J]. Genome Biology. 2005; 3: 343.
- Repp R, Hvan Ojik H, Valerius T, Groenewegen G, Wieland G, Oetzel C, et al. Phase I clinical trial of the bispecific antibody MDX-H210(anti-FcγRI anti-HER-2/neu) in combination with Filgrastim (G-CSF) for treatment of advanced breast cancer. British Journal of Cancer. 2003; 89: 2234-2243.
- Alinari L, Lapalombella R, Andritsos L, Baiocchi RA, Lin TS, Byrd JC, et al. Alemtuzumab (Campath-1H) in the treatment of chronic lymphocytic leukemia

- [J]. Nature. 2007; 26: 3644- 3553.
30. Jane C, Wilbert H, Hauger SB, Janai H, Bastian JF, Wohrley JD, et al. Infliximab treatment for refractory Kawasaki syndrome, The Journal of Pediatrics. 2005; 146: 662-667.
31. Baloch AR, Baloch AW, Sutton BJ, Zhang X. Antibody Mimetics: Promising complementary agents to animal-sourced antibodies. Critical Reviews in Biotechnology. DOI: 10.3109/07388551.2014.958431.